

Name: _____
AP Calculus AB Midterm Review

Date: _____

Midterm Exam: Wednesday, January, 24th, 12:15 – 1:45, in the cafeteria

Format:






Part I: 7 multiple choice questions with calculator (4pts each)

Part II: 9 multiple choice questions without calculator (4pts each)


Part III: 3 free response questions (12pts each)

The following sets of questions are not comprehensive; they are a mere sampling of the topics that we have covered this semester. To fully prepare for this exam, review your old exams, your notes and homework assignments.


1. The function f has the property that $f(x)$, $f'(x)$ and $f''(x)$ are negative for all real values x . Which of the following could be the graph of f ?



2. The graph of a function f is shown below. At which value of x is f continuous, but not differentiable?
- (a) a (b) b (c) c (d) d (e) e



3. The graph of $f'(x)$, the derivative of the function f , is shown below. Which of the following statements is true about f ?
- (a) f is decreasing for $-1 \leq x \leq 1$.
 (b) f is increasing for $-2 \leq x \leq 0$.
 (c) f is increasing for $1 \leq x \leq 2$.
 (d) f has a local minimum at $x = 0$.
 (e) f is not differentiable at $x = -1$ and $x = 1$.



4. $\lim_{h \rightarrow 0} \frac{(1+h)^6 - 1}{h}$ is
 (a) 0 (b) 1 (c) 6 (d) ∞ (e) nonexistent

5. $\lim_{h \rightarrow 0} \frac{\sqrt[3]{8+h} - 2}{h}$ is
 (a) 0 (b) $\frac{1}{12}$ (c) 1 (d) 192 (e) ∞

6. $\lim_{h \rightarrow 0} \frac{\ln(e+h) - 1}{h}$ is
 (a) 0 (b) $\frac{1}{e}$ (c) 1 (d) e (e) nonexistent

7. $\lim_{x \rightarrow 0} \frac{\cos x - 1}{x}$ is
 (a) -1 (b) 0 (c) 1 (d) ∞ (e) none of these

8.


x	0	$0 < x < 1$	1	$1 < x < 2$	2	$2 < x < 3$	3	$3 < x < 4$
$f(x)$	-1	Negative	0	Positive	2	Positive	0	Negative
$f'(x)$	4	Positive	0	Positive	DNE	Negative	-3	Negative
$f''(x)$	-2	Negative	0	Positive	DNE	Negative	0	Positive

Let f be a function that is continuous on the interval $[0, 4)$. The function f is twice differentiable except at $x = 2$. The function f and its derivatives have the properties indicated in the table above, where DNE indicates that the derivatives of f do not exist at $x = 2$.

(a) For $0 < x < 4$, find all values of x at which f has a relative extremum. Determine whether f has a relative maximum or a relative minimum at each of these values. Justify your answer.

(b) On the axes provided, sketch the graph of a function that has all the characteristics of f .

(Note: Use the axes provided in the pink test booklet.)



9. Consider the curve given by $y^2 = 2 + xy$




(a) Show that $\frac{dy}{dx} = \frac{y}{2y-x}$.

(b) Find all points (x, y) on the curve where the line tangent to the curve has a slope $\frac{1}{2}$.

(c) Show that there are no points (x, y) on the curve where the line tangent to the curve is horizontal.


(d) Let x and y be functions of time t that are related by the equation $y^2 = 2 + xy$. At time $t = 5$, the value of y is 3 and $\frac{dy}{dt} = 6$. Find the value of $\frac{dx}{dt}$ at time $t = 5$.

10. For which of the following does $\lim_{x \rightarrow 4} f(x)$ exist ?



- (a) I only (b) II only (c) III only (d) I and II only (e) I and III only

11. The second derivative of the function f is given by $f''(x) = x(x - a)(x - b)^2$. The graph of $f''(x)$ is shown below. For what values of x does the graph of f have a point of inflection?



- (a) 0 and a only
 (b) 0 and m only
 (c) b and j only
 (d) 0, a and b
 (e) b , j and k

Selected Calculator Active Questions

12. The position of a particle moving on the x -axis at time $t > 0$ seconds is: $x(t) = e^t - \sqrt{t}$ feet.
- (a) Find the average velocity of the particle over the interval $1 \leq t \leq 3$.
 (b) In what direction and how fast is the particle moving at $t = 1$ seconds?
 (c) When is the particle moving to the right?
 (d) Find the position of the particle when its velocity is zero.
13. Let f be the function given by $f(x) = 3e^{2x}$ and let g be the function given by $g(x) = 6x^3$. At what value of x do the graphs of f and g have parallel tangent lines?
- (a) -0.701 (b) -0.567 (c) -0.391 (d) -0.302 (e) -0.258
14. The first derivative of the function f is given by $f'(x) = \frac{\cos^2 x}{x} - \frac{1}{5}$. How many critical values does f have on the open interval $(0, 10)$?
- (a) One (b) Three (c) Four (d) Five (e) Seven

15. Which of the following is an equation of the line tangent to the graph of $f(x) = x^4 + 2x^2$ at the point where $f'(x) = 1$?
- (a) $y = 8x - 5$ (b) $y = x + 7$ (c) $y = x + 0.763$
 (d) $y = x - 0.122$ (e) $y = x - 2.146$
16. Let f be the function given by $f(x) = \cos(2x) + \ln(3x)$. What is the least value of x at which the graph of f changes concavity?
- (a) 0.56 (b) 0.93 (c) 1.18 (d) 2.38 (e) 2.44
17. If $f(x) = \sqrt[5]{x^3 - 2x}$, then $f'(\sqrt{3}) =$
- (a) 0.129 (b) 0.902 (c) 0.906 (d) 1.116 (e) 2.173

More Non-calculator Questions

18. Given: $f(x) = 5x^3 + x$. If $g(x) = f^{-1}(x)$, find $g'(6)$.
19. Find the points of discontinuity, if any: $f(x) = \begin{cases} 2x + 3 & \text{if } x \leq 4 \\ 7 + \frac{16}{x} & \text{if } x > 4 \end{cases}$
20. Find a value for the constant k , if possible, that will make the function continuous:
- (a) $f(x) = \begin{cases} 7x - 2 & \text{if } x \leq 1 \\ kx^2 & \text{if } x > 1 \end{cases}$
- (b) $f(x) = \begin{cases} kx^2 & \text{if } x \leq 2 \\ 2x + k & \text{if } x > 2 \end{cases}$
21. $\lim_{x \rightarrow +\infty} \sin\left(\frac{\pi x}{2 - 3x}\right)$
22. $\lim_{\theta \rightarrow 0} \frac{\sin 3\theta}{\theta}$

23. $\lim_{x \rightarrow 0} \frac{\tan 7x}{\sin 3x}$

24. $\lim_{\theta \rightarrow 0} \frac{\theta^2}{1 - \cos \theta}$

25. Find a nonzero value for the constant k that makes

$$f(x) = \begin{cases} \frac{\tan kx}{x} & \text{if } x < 0 \\ 3x + 2k^2 & \text{if } x \geq 0 \end{cases}$$


continuous at $x = 0$.

26. A particle moves on a line away from its initial position so that after t hours it is $s = 3t^2 + t$ miles away from its initial position.


- (a) Find the average velocity of the particle over the interval $[1, 3]$.
- (b) Find the instantaneous velocity at $t = 1$.
- (c) Find the value of t on the interval $[1, 3]$ where the instantaneous velocity is equal to the average velocity.

27. Match the graphs of the functions shown in (a) – (f) with the graphs of their derivatives in (A) – (F).


(a)




(b)




(c)




(d)




(e)




(f)




(A)




(B)




(C)




(D)



(E)



(F)



28. Given the following table of values, find the indicated derivatives:

x	$f(x)$	$f'(x)$
2	1	7
8	5	-3

(a) $g'(2)$ where $g(x) = [f(x)]^3$


(b) $h'(2)$ where $h(x) = f(x^3)$

29. Given $f(x) = \tan(4x^2)$, find $f'(x)$.


30. Given $f(x) = \sin\left(\frac{1}{x^2}\right)$, find $f'(x)$.

31. Use the graph of the equation $y = f(x)$ in the accompanying figure to find the


signs of $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ at the points A, B, and C.



32. Use the graph of $y = f''(x)$ in the accompanying figure to determine the x -coordinates of all inflection points of f . Explain your reasoning.



33. Use the graph of $y = f'(x)$ in the accompanying figure to replace the question mark with $<$, $=$, or $>$, as appropriate. Explain your reasoning.



- (a) $f(0) ? f(1)$
- (b) $f(1) ? f(2)$
- (c) $f'(0) ? 0$
- (d) $f'(1) ? 0$
- (e) $f''(0) ? 0$
- (f) $f''(2) ? 0$

For #s 34 and 35, find (a) intervals on which f is increasing, (b) intervals on which f is decreasing, (c) intervals on which f is concave up, (d) intervals on which f is concave down, and (e) x -coordinates of all points of inflection on f .

34. $f(x) = 3x^4 - 4x^3$

35. $f(x) = e^{-\frac{x^2}{2}}$

36. Given $y = \sin^{-1}\left(\frac{1}{3}x\right)$, find $\frac{dy}{dx}$.

37. Verify that the hypotheses of the Mean Value Theorem are satisfied on the given interval and find all values of c in that interval that satisfy the conclusion of the theorem for $f(x) = \sqrt{x+1}$ on the interval $[0, 3]$.

38. Given $y = \sqrt{\ln x}$, find $y'(e^5)$.

39. Given $y = e^{7x}$, find $y'(\ln 5)$.

40. Given $y = \pi^{\sin x}$, find $y'(\pi)$.

41. Find the absolute maximum and minimum values of f on the given closed interval and state where those values occur for $f(x) = \sin x - \cos x$ on the interval $[0, \pi]$.

42. Oil spilled from a ruptured oil tanker spreads in a circle whose area increases at a constant rate of 6 square miles per hour. How fast is the radius of the spill increasing when the area is 9 square miles?

43. The position of a particle moving along the line $y = 2$ is given by $x(t) = 2t^3 - 21t^2 + 60t - 50$ where t is the time in seconds, $t \geq 0$ and x is the position in feet from the point $(0, 2)$.

- (a) At what time(s) is the particle at rest?
- (b) At what time(s) is the particle moving to the right?
- (c) At what time(s) is the particle moving to the left?
- (d) What is the maximum speed of the particle on the interval $[1, 6]$?
- (e) On what interval is the velocity of the particle increasing?
- (f) What is the total distance travelled by the particle on the interval $[1, 6]$?
- (g) On what interval(s) is the speed increasing?
- (h) On what interval(s) is the speed decreasing?