

Name: \_\_\_\_\_  
AP Calc

Date: \_\_\_\_\_  
Ms. Loughran

$$f'(x) = \begin{cases} 6ax + 2b & x \leq 1 \\ 4ax^3 - 8bx - 3 & x > 1 \end{cases}$$

Do Now:

1. If the function  $f(x) = \begin{cases} 3ax^2 + 2bx + 1, & x \leq 1 \\ ax^4 - 4bx^2 - 3x, & x > 1 \end{cases}$  is differentiable for all real values of  $x$ , then  $b =$

- (A) 0      (B)  $-\frac{11}{4}$       (C)  $\frac{1}{4}$       (D)  $-\frac{7}{16}$       (E)  $-\frac{1}{4}$

continuous

$$3ax^2 + 2bx + 1 = ax^4 - 4bx^2 - 3x \quad @x=1$$

$$3a + 2b + 1 = a - 4b - 3$$

$$2a + 6b = -4$$

$$(a + 3b = -2) \quad \leftarrow -2$$

diff

$$6ax + 2b = 4ax^3 - 8bx - 3 \quad @x=1$$

$$6a + 2b = 4a - 8b - 3$$

$$2a + 10b = -3$$

$$-2a - 6b = 4$$

$$4b = 1$$

$$b = \frac{1}{4}$$

2. The position of a particle moving along the  $x$ -axis at time  $t$  is given by  $x(t) = e^{\cos 2t}$ ,  $0 \leq t \leq \pi$ . For which of the following values of  $t$  will  $x'(t) = 0$ ?

- I.  $t = 0$  ✓  
II.  $t = \frac{\pi}{2}$  ✓  
III.  $t = \pi$  ✓


- (A) I only      (B) II only      (C) I and III only      (D) I and II only      (E) I, II and III

$$x'(t) = e^{\cos 2t} \cdot -\sin 2t \cdot 2$$

$$x'(t) = -2e^{\cos 2t} \sin 2t$$

$$-2e^{\cos 2t} \sin 2t = 0$$

$$\frac{-2e^{\cos 2t} \neq 0}{\sin 2t = 0}$$



$2t = 0, \pi, 2\pi, 3\pi, \dots$   
 $t = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, \dots$  outside interval

Finishing up #12 from yesterday's packet:

$$12. f(x) = \begin{cases} ax^{-1}, & x \leq 1 \\ 12 - bx^2, & x > 1 \end{cases} \text{ at } x=1$$

$$f'(x) = \begin{cases} -ax^{-2} \text{ or } \frac{-a}{x^2} & x \leq 1 \\ -2bx & x > 1 \end{cases}$$

Cont:

$$\frac{a}{x} = 12 - bx^2 \text{ @ } x=1$$

$$a = 12 - b$$

diff

$$\frac{-a}{x^2} = -2bx \text{ @ } x=1$$

$$-a = -2b$$

$$a = 2b$$

$$12 - b = 2b$$

$$12 = 3b$$

$$4 = b$$

$$a = 12 - 4 = 8$$

# Differentiability Homework

## Homework 10-25

(1)  $y = x^5 \sec\left(\frac{1}{x}\right)$

$$y' = 5x^4 \cdot \sec\left(\frac{1}{x}\right) + \sec\left(\frac{1}{x}\right) \tan\left(\frac{1}{x}\right) \cdot \frac{-1}{x^2} \cdot x^5$$

$$y' = 5x^4 \sec\left(\frac{1}{x}\right) - x^3 \sec\left(\frac{1}{x}\right) \tan\left(\frac{1}{x}\right)$$

(2)  $y = \cos(\cos x)$

$$y' = -\sin(\cos x) \cdot -\sin x$$

$$y' = \sin(\cos x) \sin x$$

(3)  $y = x \cos 3x \quad x = \pi$

$$y' = 1 \cdot \cos 3x + x \cdot -\sin(3x) \cdot 3$$

$$y' = \cos 3x - 3x \sin(3x) \Big|_{x=\pi} \quad \cos 3\pi - 3\pi \sin 3\pi = -1$$

$$y = \pi \cos 3\pi$$

$$y = \pi(-1) = -\pi$$

(4) (4)  $y = \cot^3(\pi - \theta) \quad \frac{dy}{d\theta}$

$$y' = 3 \cot^2(\pi - \theta) \cdot -\csc^2(\pi - \theta) \cdot -1$$

$$y' = 3 \cot^2(\pi - \theta) \csc^2(\pi - \theta)$$

$$y + \pi = -(x - \pi)$$

$$y + \pi = -x + \pi$$

$$y = -x$$

(5)  $f(x) = \begin{cases} x^2 & x \leq 1 \\ \sqrt{x} & x > 1 \end{cases}$  continuous

$$f'(x) = 2x$$

$$f'(x) = \frac{1}{2}x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}}$$

not differentiable at 1

$$(6) \quad 3x^2 = ax + b \quad @ \quad x=1$$

$$3(1)^2 = a(1) + b$$

$$3 = a + b$$

$$f(x) = \begin{cases} 3x^2 & x \leq 1 \\ ax + b & x > 1 \end{cases}$$

$$f'(x) = \begin{cases} 6x & x \leq 1 \\ a & x > 1 \end{cases}$$

$$\begin{cases} a = 6 \\ b = -3 \end{cases}$$

$$6x = a$$

$$6(1) = a$$

$$6 = a$$

Since

$$3 = a + b$$

$$3 = 6 + b$$

$$b = -3$$

$$(7) \quad \frac{d^{87}}{dx^{87}} [\sin x] = -\cos x$$

$$(8) \quad \frac{d^{100}}{dx^{100}} \cos x = \cos x$$

$$d/dx = \cos x$$

$$d^2/dx^2 = -\sin x$$

$$d^3/dx^3 = -\cos x$$

$$d^4/dx^4 = \sin x$$

$$d^5/dx^5 = \cos x$$



like previous of :-

$$d/dx = -\sin x$$

$$d^2/dx^2 = -\cos x$$

$$d^3/dx^3 = \sin x$$

$$d^4/dx^4 = \cos x$$



$$(9) \quad (a) \text{ all } x$$

$$(b) \text{ all } x$$

$$(c) \quad x \neq \frac{k\pi}{2} \quad k \in \text{odd}$$

$$(d) \quad x \neq n\pi, \quad n \in \mathbb{Z}$$

$$(e) \quad x \neq \frac{\pi}{2} + n\pi \quad n \in \mathbb{Z}$$
  
 or  $\frac{\pi}{2}k \quad k \in \mathbb{Z}^{\text{odd}}$

$$(f) \quad x \neq n\pi, \quad n \in \mathbb{Z}$$

$$(g) \quad x \neq (n)\pi \quad n \in \text{odd } \mathbb{Z}$$

$$(h) \quad x \neq \frac{n\pi}{2}, \quad n \in \mathbb{Z} \quad \text{- includes } 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, \dots$$

$$(i) \text{ all } x$$

$$1 + \cos x = 0$$


$$2 - \sin x = 0$$

$$\sin x \cos x = 0$$

$$\frac{\pi}{2}, \frac{3\pi}{2} \quad \cos x = -1$$

$$\sin x = -2$$

$$\sin x = 0 \quad \cos x = 0$$



$$\sin x = \frac{1}{2}$$



(40) If  $f$  has a deriv @  $a$  then it is cont @  $a$  True

(41) If  $f$  is cont @  $x=0$ , then it has a deriv. @  $x=0$  False

$$f(x) = \begin{cases} 2x+1 & x \leq 0 \\ x^2+1 & x > 0 \end{cases}$$

(44) LHD  $2$  |  $x=0$   $2$  (B)

(45) RHD  $2x$  |  $x=0$   $0$  (C)

For 1 and 2, find  $\frac{dy}{dx}$ .

1.  $y = x^5 \sec\left(\frac{1}{x}\right)$

2.  $y = \cos(\cos x)$

3. Find an equation for the tangent line to  $y = x \cos 3x$  at  $x = \pi$ .

4. If  $y = \cot^3(\pi - \theta)$ , find  $\frac{dy}{d\theta}$ .

5. Let

$$f(x) = \begin{cases} x^2, & x \leq 1 \\ \sqrt{x}, & x > 1 \end{cases}$$

Determine whether  $f$  is differentiable at  $x = 1$ . If so, find the value of the derivative there.

6. Let

$$f(x) = \begin{cases} 3x^2, & x \leq 1 \\ ax + b, & x > 1 \end{cases}$$

Find the values of  $a$  and  $b$  so that  $f$  will be differentiable at  $x = 1$ .

7.  $\frac{d^{87}}{dx^{87}}[\sin x]$

8.  $\frac{d^{100}}{dx^{100}}[\cos x]$

9. In each part, determine where  $f$  is differentiable.

(a)  $f(x) = \sin x$  *all x*

(f)  $y = \csc x$

(i)  $f(x) = \frac{\cos x}{2 - \sin x}$  *all x*

(b)  $f(x) = \cos x$  *all x*

(g)  $f(x) = \frac{1}{1 + \cos x}$  *cos x = -1*

*2 - sin x = 0  
2 = sin x*

(c)  $f(x) = \tan x$


(h)  $f(x) = \frac{1}{\sin x \cos x}$

*$\frac{\sin x \cos x = 0}{\sin x = 0 \mid \cos x = 0}$*

(d)  $f(x) = \cot x$

(e)  $f(x) = \sec x$

p. 115 #s 40, 41, 44, 45



36.  $C(x) = x|x|$

37. Show that the function

$$f(x) = \begin{cases} 0, & -1 \leq x < 0 \\ 1, & 0 \leq x \leq 1 \end{cases}$$

is not the derivative of any function on the interval  $-1 \leq x \leq 1$ .

38. **Writing to Learn** Recall that the numerical derivative (NDER) can give meaningless values at points where a function is not differentiable. In this exercise, we consider the numerical derivatives of the functions  $1/x$  and  $1/x^2$  at  $x = 0$ .

- (a) Explain why neither function is differentiable at  $x = 0$ .
- (b) Find NDER at  $x = 0$  for each function.
- (c) By analyzing the definition of the symmetric difference quotient, explain why NDER returns wrong responses that are so different from each other for these two functions.

39. Let  $f$  be the function defined as

$$f(x) = \begin{cases} 3 - x, & x < 1 \\ ax^2 + bx, & x \geq 1 \end{cases}$$

where  $a$  and  $b$  are constants.

- (a) If the function is continuous for all  $x$ , what is the relationship between  $a$  and  $b$ ?
- (b) Find the unique values for  $a$  and  $b$  that will make  $f$  both continuous and differentiable.

### Standardized Test Questions

You may use a graphing calculator to solve the following problems.

- 40. **True or False** If  $f$  has a derivative at  $x = a$ , then  $f$  is continuous at  $x = a$ . Justify your answer.
- 41. **True or False** If  $f$  is continuous at  $x = a$ , then  $f$  has a derivative at  $x = a$ . Justify your answer.
- 42. **Multiple Choice** Which of the following is true about the graph of  $f(x) = x^{4/5}$  at  $x = 0$ ?
  - (A) It has a corner.
  - (B) It has a cusp.
  - (C) It has a vertical tangent.
  - (D) It has a discontinuity.
  - (E)  $f(0)$  does not exist.
- 43. **Multiple Choice** Let  $f(x) = \sqrt[3]{x - 1}$ . At which of the following points is  $f'(a) \neq \text{NDER}(f(x), x, a)$ ?
  - (A)  $a = 1$     (B)  $a = -1$     (C)  $a = 2$     (D)  $a = -2$
  - (E)  $a = 0$

In Exercises 44 and 45, let

$$f(x) = \begin{cases} 2x + 1, & x \leq 0 \\ x^2 + 1, & x > 0 \end{cases} \quad f'(x) = \begin{cases} 2 & x \leq 0 \\ 2x & x > 0 \end{cases}$$

- 44. **Multiple Choice** Which of the following is equal to the left-hand derivative of  $f$  at  $x = 0$ ?
  - (A)  $2x$     (B)  $2$     (C)  $0$     (D)  $-\infty$     (E)  $\infty$

45. **Multiple Choice** Which of the following is equal to the right-hand derivative of  $f$  at  $x = 0$ ?

- (A)  $2x$     (B)  $2$     (C)  $0$     (D)  $-\infty$     (E)  $\infty$

### Explorations

- 46. (a) Enter the expression " $x < 0$ " into Y1 of your calculator using " $<$ " from the TEST menu. Graph Y1 in DOT MODE in the window  $[-4.7, 4.7]$  by  $[-3.1, 3.1]$ .
  - (b) Describe the graph in part (a).
  - (c) Enter the expression " $x \geq 0$ " into Y1 of your calculator using " $\geq$ " from the TEST menu. Graph Y1 in DOT MODE in the window  $[-4.7, 4.7]$  by  $[-3.1, 3.1]$ .
  - (d) Describe the graph in part (c).

47. **Graphing Piecewise Functions on a Calculator** Let

$$f(x) = \begin{cases} x^2, & x \leq 0 \\ 2x, & x > 0 \end{cases}$$

- (a) Enter the expression " $(X^2)(X \leq 0) + (2X)(X > 0)$ " into Y1 of your calculator and draw its graph in the window  $[-4.7, 4.7]$  by  $[-3, 5]$ .
- (b) Explain why the values of Y1 and  $f(x)$  are the same.
- (c) Enter the numerical derivative of Y1 into Y2 of your calculator and draw its graph in the same window. Turn off the graph of Y1.
- (d) Use TRACE to calculate  $\text{NDER}(Y1, x, -0.1)$ ,  $\text{NDER}(Y1, x, 0)$ , and  $\text{NDER}(Y1, x, 0.1)$ . Compare with Section 3.1, Example 6.

### Extending the Ideas

48. **Oscillation** There is another way that a function might fail to be differentiable, and that is by *oscillation*. Let

$$f(x) = \begin{cases} x \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

- (a) Show that  $f$  is continuous at  $x = 0$ .
- (b) Show that

$$\frac{f(0 + h) - f(0)}{h} = \sin \frac{1}{h}$$

- (c) Explain why

$$\lim_{h \rightarrow 0} \frac{f(0 + h) - f(0)}{h}$$

does not exist.

- (d) Does  $f$  have either a left-hand or right-hand derivative at  $x = 0$ ?
- (e) Now consider the function

$$g(x) = \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Use the definition of the derivative to show that  $g$  is differentiable at  $x = 0$  and that  $g'(0) = 0$ .