

36. $C(x) = x|x|$

37. Show that the function

$$f(x) = \begin{cases} 0, & -1 \leq x < 0 \\ 1, & 0 \leq x \leq 1 \end{cases}$$

is not the derivative of any function on the interval $-1 \leq x \leq 1$.

38. **Writing to Learn** Recall that the numerical derivative (NDER) can give meaningless values at points where a function is not differentiable. In this exercise, we consider the numerical derivatives of the functions $1/x$ and $1/x^2$ at $x = 0$.
- (a) Explain why neither function is differentiable at $x = 0$.
- (b) Find NDER at $x = 0$ for each function.
- (c) By analyzing the definition of the symmetric difference quotient, explain why NDER returns wrong responses that are so different from each other for these two functions.

39. Let f be the function defined as

$$f(x) = \begin{cases} 3 - x, & x < 1 \\ ax^2 + bx, & x \geq 1 \end{cases}$$

where a and b are constants.

- (a) If the function is continuous for all x , what is the relationship between a and b ?
- (b) Find the unique values for a and b that will make f both continuous and differentiable.

Standardized Test Questions

You may use a graphing calculator to solve the following problems.

40. **True or False** If f has a derivative at $x = a$, then f is continuous at $x = a$. Justify your answer.
41. **True or False** If f is continuous at $x = a$, then f has a derivative at $x = a$. Justify your answer.
42. **Multiple Choice** Which of the following is true about the graph of $f(x) = x^{4/5}$ at $x = 0$?
- (A) It has a corner.
 (B) It has a cusp.
 (C) It has a vertical tangent.
 (D) It has a discontinuity.
 (E) $f(0)$ does not exist.
43. **Multiple Choice** Let $f(x) = \sqrt[3]{x-1}$. At which of the following points is $f'(a) \neq \text{NDER}(f(x), x, a)$?
- (A) $a = 1$ (B) $a = -1$ (C) $a = 2$ (D) $a = -2$
 (E) $a = 0$

In Exercises 44 and 45, let

$$f(x) = \begin{cases} 2x + 1, & x \leq 0 \\ x^2 + 1, & x > 0 \end{cases}$$

44. **Multiple Choice** Which of the following is equal to the left-hand derivative of f at $x = 0$?
- (A) $2x$ (B) 2 (C) 0 (D) $-\infty$ (E) ∞

45. **Multiple Choice** Which of the following is equal to the right-hand derivative of f at $x = 0$?

(A) $2x$ (B) 2 (C) 0 (D) $-\infty$ (E) ∞

Explorations

46. (a) Enter the expression " $x < 0$ " into Y1 of your calculator using "<" from the TEST menu. Graph Y1 in DOT MODE in the window $[-4.7, 4.7]$ by $[-3.1, 3.1]$.
- (b) Describe the graph in part (a).
- (c) Enter the expression " $x \geq 0$ " into Y1 of your calculator using " \geq " from the TEST menu. Graph Y1 in DOT MODE in the window $[-4.7, 4.7]$ by $[-3.1, 3.1]$.
- (d) Describe the graph in part (c).

47. **Graphing Piecewise Functions on a Calculator** Let

$$f(x) = \begin{cases} x^2, & x \leq 0 \\ 2x, & x > 0 \end{cases}$$

- (a) Enter the expression " $(X^2)(X \leq 0) + (2X)(X > 0)$ " into Y1 of your calculator and draw its graph in the window $[-4.7, 4.7]$ by $[-3, 5]$.
- (b) Explain why the values of Y1 and $f(x)$ are the same.
- (c) Enter the numerical derivative of Y1 into Y2 of your calculator and draw its graph in the same window. Turn off the graph of Y1.
- (d) Use TRACE to calculate $\text{NDER}(Y1, x, -0.1)$, $\text{NDER}(Y1, x, 0)$, and $\text{NDER}(Y1, x, 0.1)$. Compare with Section 3.1, Example 6.

Extending the Ideas

48. **Oscillation** There is another way that a function might fail to be differentiable, and that is by *oscillation*. Let

$$f(x) = \begin{cases} x \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

- (a) Show that f is continuous at $x = 0$.
- (b) Show that

$$\frac{f(0+h) - f(0)}{h} = \sin \frac{1}{h}$$

- (c) Explain why

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$$

does not exist.

- (d) Does f have either a left-hand or right-hand derivative at $x = 0$?
- (e) Now consider the function

$$g(x) = \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Use the definition of the derivative to show that g is differentiable at $x = 0$ and that $g'(0) = 0$.